REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Artington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE	3. REPORT TYPE A	ND DATES COVERED	
	12/15/97		/01/93 to 07/31/97	
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS	
Method Development for Un Enrichment Experiment	G N00014-94-1-0125			
6. AUTHOR(S)				
Dr. Kenneth Johnson Dr. Kenneth Coale				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			8. PERFORMING ORGANIZATION REPORT NUMBER	
San Jose State University Foundation (Moss Landing Marine Labs) P.O. Box 720130				
San Jose CA 951720-0130			21-1509-0494	
9. SPONSORING/MONITORING AGENCY	NAME(S) AND ADDRESS(ES)		10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
Office of Naval Research				
Seattle Regional Office 1107 NE 45th Street, Suit				
Seattle WA 98105-4631				
44 CURRIENTARY MOTES				
11. SUPPLEMENTARY NOTES				
			T	
12a. DISTRIBUTION/AVAILABILITY STATEMENT			12b. DISTRIBUTION CODE	
DISTRO	BUTION STATEMENT E			
	ved tox public released			
12 ARCTRACT (Maximum 200 words)				

13. ABSTRACT (Maximum 200 words)

The long-term goals of this work were to test the hypothesis that metals, especially iron, regulate rates of primary production in High Nutrient, Low Chlorophyll (HNLC) areas of the ocean. Two open ocean, iron enrichments were conducted in the eastern equatorial Pacific. Four nM iron was added in a single addition during the first experiment (Oct. 1993) and in three separate additions of 2, 1 and 1 nM in the second (May 1995). Ther: was a strong response of the ecosystem with increases in primary production rates and plankton biomass. Large changes in bio-optical properties were also observed. The second experiment, in which higher iron concentrations were sustained for 1 week, produced a 20-fold increase in chlorophyll - a proxy for photoplankton biomass. The results of the experiments have been reported in some 20 journal articles.

ŀ	14. SUBJECT TERMS	15. NUMBER OF PAGES		
	Ocean Cemistry, Ocean	16. PRICE CODE		
	17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
	unclassified	unclassified	unclassified	200 (Day 2 80)

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std 239-18 298-102

Kenneth S. Johnson & Kenneth H. Coale

METHOD DEVELOPMENT FOR AN UNENCLOSED IRON FERTILIZATION EXPERIMENT N00014-94-1-0125

Moss Landing Marine Laboratories
PO Box 450
Moss Landing, CA 95039
johnson@mlml.calstate.edu, coale@mlml.calstate.edu
408 755 8657 tel
408 753 2826 fax

The long-term goals of this work were to test the hypothesis that metals, especially iron, regulate rates of primary production in High Nutrient, Low Chlorophyll (HNLC) areas of the ocean. An important corollary of this hypothesis is that it might be feasible to regulate productivity in small patches in the open ocean by addition of small amounts of iron. The ability to produce "mini- blooms" would be an extremely valuable experimental tool for the study of ocean biogeochemistry, optical oceanography and remote sensing.

The primary objectives of the work were to perform the first mesoscale (64 km²) iron enrichment experiments in the open ocean. The community response to these iron additions in HNLC waters was then determined. The first enrichment experiment was conducted in October/November, 1993 in the HNLC waters south of the Galapagos Islands at 90°W. The second was performed further west of the Islands (103°W). Iron (400 kg) was added with sulfur hexafluoride (an inert tracer) to 64 km² patches in both experiments. All of the iron was added in a single addition, bringing the mixed layer concentration to 4 nM in the first experiment. The iron was added in 3 separate additions over a 1 week duration in the second experiment to concentrations of 2, 1 and 1 nM.

The first iron enrichment experiment was a major success. Addition of iron to a 64 km² patch produced increases in chlorophyll concentration and primary production rates that were 3-fold elevated over background values (Martin et al., 1994). We were able to track this patch for 9 days before returning to port. These results demonstrate that iron does regulate biological rates in HNLC areas of the ocean. They also clearly indicate the feasibility of performing controlled experiments on open ocean ecosystems for significant periods of time. It is clear, however, that the ecosystem in the fertilized patch did not behave similarly to that observed in containers on board ship where all of the available nitrate is depleted when iron is added and chlorophyll may increase 30-fold or more. Three hypotheses might explain the lack of nitrate depletion observed in IronEx I: 1) another trace nutrient such as zinc was depleted, 2) grazing in the patch limited the growth of phytoplankton or 3) the iron was lost from the patch so rapidly that little was taken up by the phytoplankton and it rapidly became limiting again.

To test these hypotheses, iron was repeatedly added to the enriched patch during IronEx II in 1995. This patch was tracked over a distance of 1100 km during an 18 day span. Chlorophyll increased more than 20-fold during this experiment (Coale et al., 1996). At the same time, about one half of the available nitrate was consumed and the partial pressure of carbon dioxide was drawn down as much as 100 uatm (Cooper et al., 1996). Large amounts of

19971222 028

DITIC QUALITY INSPECTED 4

dimethyl sulfide were produced, as well (Turner et al., 1996). This result clearly eliminates the possibility that grazers or a second micronutrient limited the patch.

The results of the 1993 cruise have been summarized in 3 papers in Nature (Martin et al., Nature, 371, 123, 1994; Kolber et al., Nature, 371, 145, 1994; Watson et al., 371, 143, 1994). The results from all of the investigators in the 1993 cruise are in press at Deep-Sea Research II as a special issue of the journal.

The first results from the 1995 IronEx II cruise were published as 4 papers in Nature (Coale et al., Nature, 383,495-501, 1996; Behrenfeld et al., Nature, 383, 509, 1996; Cooper et al., Nature, 383, 511, 1996; Turner et al., Nature 383, 513, 1996). Several additional papers have been published by other investigators participating in the ONR funded expedition. This includes a paper by Kudela et al. (1996) on the optical properties of the iron enriched patch. Additional papers are in preparation including several from our group.

The two IronEx experiments have broad implications for how oceanography is performed. They demonstrate that we can move from an observational science into an era of controlled experimentation that will rapidly advance our understanding of chemistry, biology and optics. For example, deliberate experiments can now be conducted to assess the role of ecosystems in producing Colored Dissolved Organic Matter (CDOM) or biogenic aerosols such as DMS (e.g. Turner et al., 1996). The partitioning of natural isotopes in these experimental systems can be used to ground truth the processes that lead to biomarker formation and fractionation, such as alkenone temperature index.

The IronEx experiments have strong implications for the role of oceanic biology in regulating the climate of the earth. We are now working with NSF to plan a 3rd IronEx experiment in the Southern Ocean. Iron regulation of plankton growth in this region must be demonstrated if the iron-climate linkage is to be confirmed. In addition, we are working with the Monterey Bay Aquarium Research Institute to examine the potential for iron regulation of primary production and optics in the coastal zone.

METHOD DEVELOPMENT FOR AN UNENCLOSED IRON FERTILIZATION EXPERIMENT N00014-94-1-0125

Moss Landing Marine Laboratories PO Box 450 Mcss Landing, CA 95039 johnson@mlml.calstate.edu 408 755 8657 tel 408 753 2826 fax

A. PAPERS From MLML ACKNOWLEDGING N00014-94-1-0125

- Johnson, K. S., K. H. Coale, V. A. Elrod, and N. W. Tindale. 1994. Iron Photochemistry in seawater from the Equatorial Pacific. Marine Chemistry 46, 319-334.
- Martin, J. H., K. H. Coale, K. S. Johnson, S. E. Fitzwater, R. M. Gordon, S. J. Tanner, C. N. Hunter, V. A. Elrod, J. L. Nowicki, T. L. Coley, R. T. Barber, S. Lindley, A. J. Watson, K. Van Scoy, C. S. Law, M. I. Liddicoat, R. Ling, T. Stanton, J. Stockel, C. Collins, A. Anderson, R. Bidigare, M. Ondrusek, M. Latasa, F. J. Millero, K. Lee, W. Yao, J. Z. Zhang, G. Friederich, C. Sakamoto, F. Chavez, K. Buck, Z. Kolber, R. Greene, P. Falkowski, S. W. Chisholm, F. Hoge, R. Swift, J. Yungel, S. Turner, P. Nightingale, A. Hatton, P. Liss, N. W. Tindale. 1994. Testing the Iron Hypothesis in Ecosystems of the Equatorial Pacific Ocean. Nature, 371, 123-129.
- Kolber, Z., R. T. Barber, S. W. Chisholm, K. H. Coale, R. M. Greene, K. S. Johnson, S. Lindley, and P. G. Falkowski. 1994. Iron limits photosynthetic energy conversion efficiency in the equatorial Pacific Ocean. Nature, 371, 174-179.
- Johnson, K. S., and H. W. Jannasch. 1994. Analytical chemistry under the sea surface: monitoring ocean chemistry in situ. Naval Research Reviews, XLVI, #3, 4-12.
- Watson, A. J., C. S. Law, K. A. Van Scoy, F. J. Millero, W. Yao, G. E. Friederich, M. I. Liddicoat, R. H. Wanninkhof, R. T. Barber and K. H. Coale. 1994. Minimal effect of iron fertilization on sea-surface carbon dioxide concentrations. Nature, 371, 143-145.
- Coale, K. H., S. E. Fitzwater, R. M. Gordon, K. S. Johnson and R. T. Barber. 1996. Control of community growth and export production by upwelled iron the equatorial Pacific Ocean. Nature, 379, 621-624.
- Coale, K. H., K. S. Johnson, S. E. Fitzwater, R. M. Gordon, S. Tanner, F. P. Chavez, L. Ferioli, C. Sakamoto, P. Rogers, F. Millero, P. Steinberg, P. Nightingale, D. Cooper, W. P. Cochlan, M. R. Landry, J. Constantinou, G. Rollwagen, A. Trasvina, and R.

- Kudela. 1996. A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the equatorial Pacific Ocean. Nature, 383, 495-501.
- Gordon, R. M., K. H. Coale and K. S. Johnson. 1997. Iron distributions in the Equatorial Pacific: implications for new production. Limnology and Oceanography 42, 419-431.
- Fitzwater, S. E., K. H. Coale, R. M. Gordon, K. S. Johnson and M. E. Ondrusek. 1996. Iron deficiency and phytoplankton growth in the Equatorial Pacific. Deep-Sea Research II, 43 995-1015.
- Johnson, K. S., R. M. Gordon, and K. H. Coale. 1997. What controls dissolved iron in the world ocean? Marine Chemistry 57, 137-161.
- Johnson, K. S., R. M. Gordon, and K. H. Coale. 1997. What controls dissolved iron in the world ocean: Author's closing comments. Marine Chemistry, 57, 181-186.
- Gordon, R. M., K. S. Johnson and K. H. Coale. In press. Behavior of iron and other trace elements in IronEx I and PlumEx. Deep-Sea Research.
- Coale, K. H., S. Fitzwater, S. Blain, T. Coley, C. Hunter, K. Johnson, T. Stanton and A. Watson. In press. The design and implementation of a mesoscale iron enrichment experiment in the equatorial Pacific. Deep-Sea Research.
- Johnson, K. S., K. H. Coale, s. Tanner, C. Hunter, R. M. Gordon and V. Elrod. In preparation. The iron hypothesis in the equatorial Pacific: The role of iron in regulating phytoplankton carbon production during IronEx I and II. Limnology and Oceanography.

B. PAPERS BY OTHER RESEARCH GROUPS STEMMING FROM THE IRONEX EXPERIMENTS:

- Behrenfeld, M. J., A. J. Bale, Z. S. Kolber, J. Aiken, and P. G. Falkowski. 1996. Confirmation of iron limitation of phytoplankton photosynthesis in the equatorial Pacific Ocean. Nature, 383, 508-511.
- Copper, D. J., A. J. Watson, and P. D. Nightingale. 1996. Large decrease in ocean-surface CO₂ fugacity in response to *in situ* iron fertilization. Nature, 383, 511-513.
- Turner, S. M., P. D. Nightingale, L. J. Spokes, M. I. Liddicoat, and P. S. Liss. 1996. Increased dimethyl sulphide concentrations in sea water from *in situ* iron enrichment. Nature, 383, 513-517.
- Kudela, R. M. and F. P. Chavez. 1996. Bio-optical properties in relation to an algal bloom caused by iron enrichement in the Equatorial Pacific. Geophysical Research Letters, 23, 3751-3754.

- Hatton, A. D., S. M. Turner, G. Malin and P. S. Liss. (1997). Biogenic sulfur compounds of atmospheric significance in the Galapagos Plume. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).
- Hoge, F. E., C. W. Wright, R. N. Swift, J. K. Yungel, R. E. Berry and R. Mitchell. (1997). Fluorescence signatures of an iron-enriched phytoplankton community in the Eastern Equatorial Pacific Ocean. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).
- Hoge, F. E., C. W. Wright, R. N. Swift, J. K. Yungel, R. E. Berry and R. Mitchell. (1997). Airborne bio-optics survey of the Galapagos islands margins. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).
- Law, C. S., A. J. Watson, M. I. Liddicoat and T. Stanton. (1997). Sulphur hexafluoride as a tracer of biogeochemical and physical processes in an open-ocean iron fertilization experiment. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).
- Lindley. S., and R. T. Barber. (1997). Phytoplankton response to natural and experimental iron addition. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).
- Millero, F. J., W. Yao, K. Lee, J-Z Zhang and D. M. Campbell. (1997). Carbonate system in the waters near the Galapagos Islands. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).
- Sakamoto, C. M., F. J. Millero, W. Yao, G. E. Friederich and F. P. Chavez. (1997). Surface seawater distributions of inorganic carbon and nutrients around the Galapagos Islands: Results from the PlumEx experiment using automated chemical mapping. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).
- Stanton, T. P, C. S. Law and A. J. Watson. (1997). Physical evolution of the IronEx I open ocean tracer patch. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).
- Steger, J. M., C. A. Collins and P. C. Chu. (1997). Circulation in the Archipielago de Colon (Galapagos Islands) November, 1993. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).
- Turner, S. M., G. Malin, K. H. Coale, M. E. Ondresek, C. S. Law and P. S. Liss. (1997). Iron fertilisation in the ocean: Effects on dimethylsulphonioproprionate and dimethyl sulphide production. J. Martin Special Issue, Deep-Sea Research vol 2, (In Press).

C. GRADUATE STUDENTS SUPPORTED

2 female - Ms. Heidi Zamzow, Ms. Laurie Ferioli 1 male - Mr. Peter von Langen

D. HONORS/AWARDS

K. S. Johnson, Sverdrup Lecturer, American Geophysical Union, 1994 - New technologies for ocean experimentation.